

Longwood University

## Digital Commons @ Longwood University

---

Fall Showcase for Research and Creative Inquiry

Office of Student Research

---

Fall 11-15-2021

### Non-local approximation Properties

Kira Pierce

Follow this and additional works at: [https://digitalcommons.longwood.edu/rci\\_fall](https://digitalcommons.longwood.edu/rci_fall)



Part of the [Computer Sciences Commons](#), and the [Mathematics Commons](#)


---

#### Recommended Citation

Pierce, Kira, "Non-local approximation Properties" (2021). *Fall Showcase for Research and Creative Inquiry*. 154.

[https://digitalcommons.longwood.edu/rci\\_fall/154](https://digitalcommons.longwood.edu/rci_fall/154)

This Poster is brought to you for free and open access by the Office of Student Research at Digital Commons @ Longwood University. It has been accepted for inclusion in Fall Showcase for Research and Creative Inquiry by an authorized administrator of Digital Commons @ Longwood University. For more information, please contact [hamiltonma@longwood.edu](mailto:hamiltonma@longwood.edu), [alwinehd@longwood.edu](mailto:alwinehd@longwood.edu).



# Non-local approximation properties of $\varphi(x) = x^{-1} \ln(1 + x^2)$

Kira Pierce  
Longwood University



<https://www.overleaf.com/project/60dbccd558b7874110a58a3e>

## Introduction

This project concerns approximation properties of the set

$$S(\varphi, X) := \left\{ \sum_{j=1}^N a_j \varphi(x - x_j) : a_j \in \mathbb{R}, x_j \in X \right\},$$

where  $X$  is a scattered sequence and  $\varphi(x) = x^{-1} \ln(1 + x^2)$ . Similar approximation sets are commonly used in interpolation problems and are especially helpful due to their Fourier representation. For our work, we will work to prove the following theorem.

### Main Result

Suppose  $f \in C[a, b]$ . For any  $\varepsilon > 0$ , there exists  $s \in S$ , such that

$$\|f - s\|_{L_\infty} < \varepsilon.$$

We begin with the Taylor Series for

$$\varphi(x - y) = (x - y)^{-1} \ln(1 + (x - y)^2)$$

which yields

$$\varphi(x - y) := \ln|y| \sum_{j=1}^{\infty} \frac{A_j(x)}{y^j} + \sum_{k=2}^{\infty} \frac{B_k(x)}{y^k}.$$

for some polynomials  $A_j(x)$  and  $B_k(x)$ . Using methods from linear algebra, we then collect  $A_j$ . Our interest was spurred by approximation theoretic results, namely those found in [1], [2], and [3].

## Cauchy Product

The Cauchy Product is the blending of two power series. Let

$$\sum_{n=0}^{\infty} a_n \text{ and } \sum_{n=0}^{\infty} b_n$$

be two series. The Cauchy Product of these two series is defined as the sum

$$\sum_{n=1}^{\infty} a_n \text{ where } c_n = \sum_{k=0}^n a_k b_{n-k} \text{ for all } n \in 0, 1, 2, \dots$$

## Vandermonde Matrix

The Vandermonde Matrix is a matrix in which each element increases in a geometric pattern by row or column.

$$V = \begin{bmatrix} 1 & x_1 & x_1^2 & \dots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \dots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_m & x_m^2 & \dots & x_m^{n-1} \end{bmatrix}$$

The determinant of a square Vandermonde matrix can be expressed as

$$\det(V) = \prod_{1 \leq i < j \leq n} (x_j - x_i).$$

## $\varphi(x) = x^{-1} \ln(1 + x^2)$

$$\begin{aligned} \frac{\partial}{\partial y} (\varphi(x - y)) &= \frac{-2(x - y)}{1 + (x - y)^2} \\ &= \sum_{n=0}^{\infty} \frac{2A_{n-1}(x)}{y^n} + \sum_{k=2}^{\infty} \frac{-2xA_{k-2}(x)}{y^k} \\ &= \frac{2A_0(x)}{y} + \sum_{k=2}^{\infty} \frac{2A_{k-1}(x) - 2xA_{k-2}(x)}{y^k} \end{aligned}$$

And from [1] we know that  $A_n(x) = (n + 1)x^n + \text{lower order terms}$

Note:  $A_0 = 1$  So we get that:

$$\frac{\partial}{\partial y} (\varphi(x - y)) = \frac{2}{y} + \sum_{k=2}^{\infty} \frac{B_k(x)}{y^k}$$

We can then write the sum of the series

$$\sum_{j=1}^{\infty} \frac{A_j(x)}{y^j} + \sum_{k=2}^{\infty} \frac{B_k(x)}{y^k}$$

With

$$A_j = \ln|y| \sum_{j=1}^{\infty} \frac{-2x^{j-1}}{y^j} \text{ and } B_k = \left[ \sum_{j=1}^{\infty} \frac{x^{j-1}}{y^j} \right] \left[ \sum_{k=1}^{\infty} \left( \frac{B_{k+1}(x)}{k} \right) \frac{1}{y^k} \right]$$

Simplifying  $A_j$ , we get that

$$A_j = -2x^{j-1}, j \geq 1.$$

The Cauchy Product is used to simplify  $B_k$ .

$$\sum_{k=1}^{\infty} \sum_{j=1}^{\infty} \frac{x^{j-1} B_{k+1}(x)}{k} \frac{1}{y^{j+k}}.$$

When  $m = j + k$  and  $j = m - k$ , we then use substitution and get

$$\sum_{m=2}^{\infty} \left( \sum_{k=1}^{m-1} \frac{x^{m-k-1} B_{k+1}(x)}{k} \right) \frac{1}{y^m}$$

where  $2 \leq m \leq \infty$  and  $1 \leq k \leq m - 1$

So,

$$C_m(x) = \sum_{k=1}^{m-1} \frac{x^{m-k-1} 2x^k}{k} = \left( \sum_{k=1}^{m-1} \frac{2}{k} \right) x^{m-1}$$

## Approximations and Differences in Functions

4th degree approximation and difference in polynomials.




Figure 1.  $\varphi(x)$  4x4 Approximation






Figure 2.  $\varphi(x)$  4x4 Difference

## Examples


### $e^{1/2x} + \cos x$



### $e^{3x} \sin 2x$



### $e^x \sin(2 \tan x)$



## Conclusion and Further Work

As shown, we know that our polynomials of  $A_j(x)$  form a basis for  $\Pi$ , therefore  $\varphi(x) = x^{-1} \ln(1 + x^2)$  can be used for its approximating properties. Furthermore, it is true that the same approximation scheme is applicable for the arbitrary values of  $q$  and  $r$  in

$$(1 + x^q)^r$$

where  $q$  is any natural number and  $r$  is any real number that is not a natural number.

## Acknowledgments and References

- Dr. Jeff Ledford for his patient guidance, enthusiastic encouragement and useful critiques of this research work.
- Brock Erwin and Calvin Foster for them working alongside me during the term, contributing insightful comments, and assisting me in completing this work.
- Office of Student Research for supporting this research and granting me this opportunity.

- J.Ledford, Approximating continuous functions with scattered translates of the Poisson kernel. Missouri J. Math. Sci. **26** (2014), no. 1, 64-69.
- J.Ledford, On the density of scattered translates of the general multiquadratic in  $C([a,b])$ . New York J. Math. **20** (2014), 145-151.
- M.J.D. Powell, Univariate multiquadric interpolation: Reproduction of linear polynomials, in *Multivariate Approximation and Interpolation (Duisberg 1989)*, Internat. Ser. Numer. Math. **94**, 227-240, Birkhäuser, Basel, 1990.